

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

7,000

Open access books available

186,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.  
For more information visit [www.intechopen.com](http://www.intechopen.com)



Chapter

# Magnetic Resonance-Guided Focused Ultrasound in the Treatment of Colorectal Cancer Liver Metastases

*Ryan Holman, Orane Lorton, Pauline C. Guillemin, Andrea Peloso, Alexis Ricoeur and Rares Salomir*

## Abstract

Liver metastases often result secondary to colorectal cancer and curative prognosis is poor. Magnetic resonance high intensity focused ultrasound is a burgeoning technique with the potential to provide a new image-guidance modality for focused ultrasound ablation of both primary and secondary liver tumors. This is particularly important for colorectal liver metastases cases ineligible for surgical resection, as chemotherapy can often be ineffective at bridging the patient for surgery, and liver transplant has generally been inadequate. At least one system for focused ultrasound ablation of primary and secondary tumors has previously been approved in the European Union, under ultrasound guidance. Magnetic resonance guidance offers many benefits, such as: integration with pre-existing imaging systems, real-time temperature mapping, and ability to assess treatment with MRI during the procedure. This chapter reviews the main aspects in treatment of this disease using this new therapy, including: focused ultrasound physics, magnetic resonance physics, magnetic resonance sequences and protocols in liver imaging, protocols and sequences in magnetic resonance thermometry, standard treatment options and limitations, relevant ongoing clinical trials, previous pilot studies, and outlooks for potential translation of this image-guidance modality as a novel ablative therapy for colorectal liver metastases.

**Keywords:** interventional radiology, focused ultrasound, liver cancer, thermal ablation, colorectal liver metastases

## 1. Introduction

Open surgery generally offers the best long-term survival rates for colorectal liver metastases (CRLM); with minimally invasive techniques becoming more common [1]. Magnetic resonance guided high intensity focused ultrasound (MRgHIFU) is noninvasive and non-ionizing, allowing for reduced treatment morbidity. At least one system for ultrasound guided focused ultrasound (USgFUS) ablation has been approved within the European Union for primary and secondary hepatic tumors [2].

Although, liver metastases are more common than primary liver tumors, most focused ultrasound studies report outcomes for primary hepatocellular carcinoma (HCC). The use of MRgHIFU for both primary and secondary hepatobiliary tumors is still awaiting certification and has not yet been reported in randomized controlled trials for CRLM or HCC [3]. Discussed here are the focused ultrasound (FUS) physics, the principles of MRI for liver metastases, analysis of the standard treatment approaches for CRLM, and previous studies involving ablation of liver tumors with USgFUS and MRgHIFU.

In 2019, cancer was reported to be the second leading cause of death, globally; amounting to approximately 1 in 6 deaths, worldwide [4]. The primary cause was due to exogenous factors resulting in genetic mutations and amounts to about 90% of reported cases [5]. P53 mutations in tumor suppressor genes are estimated in about 50% of cancers and RAS gene mutations of proto-oncogenes are estimated in about 30% of cancers. Tobacco use is thought to account for the majority of all cancer deaths. This is followed by high body mass index, alcohol use, and malnutrition [5].

HCC is the most common primary liver tumor type. There were approximately 906,000 new primary liver cancer cases in 2020, of which 75–85% were HCC, arriving at approximately 679,500–770,100 new HCC cases [6]. The most prevalent underlying conditions for HCC are Hepatitis-B virus, Hepatitis-C virus, and liver cirrhosis [7, 8]. Primary liver tumor treatment depends on history and staging. If HCC results from decompensated liver cirrhosis, surgical resection is not recommended. These patients do have the option of total liver transplant with 5-year survival rates of about 60–70%. Curative treatment options for late-stage diagnosis or recurrence is rare [8–10]. For HCC, the 10-year survival rate after surgical resection is approximately 25% [11]. However, liver transplant often offers much better outcomes than surgery for HCC. With liver transplant for patients meeting the Milan criteria, 5-year survival rates are near 70%, with less than 10% recurrence rates [11–13]. Liver transplant for HCC constitutes about 25% of liver transplants in the USA and about 40% of liver transplants in Europe [14].

CRLM is the most common form of secondary liver tumor [15]. CRLM occurs in about one-third to one-half of adult CRC cases and the liver metastases is the cause of death in about two-thirds of these patients [16]. In 2020, there were approximately 1.9 million new cases of CRC, of which it might be expected that 633,333–950,000 developed liver metastases [6]. Diagnostic radiologists have listed secondary liver tumor sites at 18–40 times more frequent than primary liver tumors, as the condition often presents with multiple metastases [17]. Historically, CRLM was deemed incurable with untreated 5-year survival rates of less than 2% [18]. Survival rates of patients with distant secondary metastatic tumors can be improved with surgical treatment and systemic chemotherapy [8, 19, 20]. Pediatric liver metastasis is more often secondary to Wilms' tumors or neuroblastomas rather than CRC [21, 22].

About 15% of adult patients exhibit liver metastasis at initial CRC diagnosis [23] and about 70% develop CRLM [2]. Approximately 60% of CRC deaths result from liver metastases [23, 24]. The standard treatment for CRLM is liver resection and is largely considered the best option for long-term curative potential [1, 8, 25, 26], with about a 40% survival rate after 5 years [2, 8, 23, 27], about a 24% survival rate after 10 years [2, 28], about a 20% cure rate [28], and a median survival rate of approximately 30 months [29]. However, for both HCC and CRLM, surgical eligibility is only 20–25% [2]. Liver transplant for CRLM has given good results in recent clinical trials when using tighter inclusion criteria and molecular profiling [30–32], although has historically given dismal survival rates, with high incidence of recurrence, survival rates only marginally better than systemic chemotherapy, and is not a primary option in

standard treatment algorithms [1, 33]. Hence, CRLM has an additional treatment difficulty, compared to HCC, because liver transplant does not generally provide long term survival.

## 2. Focused ultrasound principles

FUS surgery was first reported in 1942 after being applied to cat and dog brain tissue [34, 35] and more elaborate neurological studies later followed [36, 37]. MRgHIFU integrates a FUS transducer into a MRI system with near real-time imaging feedback; capable of temporal resolution less than 1.0 seconds, in-plane resolution less than 1.0 mm, and temperature resolution less than 1.0°C [34]. Thermal tissue ablation results from rapid temperature change of greater than 55°C during heating or –20 to –50°C during cooling [38]. Adequate ablation for coagulative necrosis requires about 10 seconds, with intermittent cooling periods to avoid skin burning [34]. More recent developments enable various feedback methods to regulate temperature, optimize speed, and automate the scanning procedure [39].

FUS works via constructive wave interference. The waves are generated by powering piezoelectric elements with an alternating current [40]. Most modern transducers are phase-array types, composed of hundreds of elements that can be individually controlled, each emitting a low amplitude ultrasonic wave at the focus [40]. Each wave is low enough in amplitude to pass through the tissue without causing significant heating, interfering constructively at the focus. The phase lag of each transducer element is adjusted so the waves are in-phase at the focal region, capable of performing beam steering and refocusing phase aberrations from bone or tissue inhomogeneities. When the waves form a large amplitude oscillation, the heating increases substantially and allows ablation and coagulative necrosis. The wave amplitude and frequency can be controlled by the operator as well as other factors like position, applied power, and pulse modulation. Lower frequencies are better for deep sites like transcranial applications, while high frequencies are used for surface sites [41].

Tissue has an inherent property to absorb ultrasonic energy. The acoustic absorption coefficient measures a tissue's ability to absorb ultrasound. In tissue at 1 MHz, the beam attenuates to about 50% at a depth of about 7 cm [38]. Beam reflection is significant at interfaces with large differences in acoustic absorption coefficient, causing high amounts of reflection at tissue-gas interfaces and tissue-bone interfaces [38]. At large FUS powers, strong rarefactional pressures exist. If this is coupled with lower frequency ultrasound waves, the conditions are favorable to induce tissue nucleation [34, 42]. This results in cavitation heating that can cause detrimental tissue damage or be utilized in techniques like lithotripsy [43, 44] and histotripsy [45, 46]. Low temperature therapies expose cells to about 43–45°C for long time periods. High temperature thermal therapy uses temperature between 50°C and 80°C for short time periods to ablate tissue, cause coagulation, and induce necrosis [47]. The tissue damage is estimated by the equivalent number of thermal doses at 43°C, with necrosis induced after about 240 min at 43°C [48, 49].

## 3. MRI principles

MRI is based on the concept of nuclear magnetic resonance. Atomic nuclei with an odd number of protons or neutrons exhibit a net spin entailing a charge circulation that forms an individual magnetic field surrounding the atom, giving the protons a

magnetic dipole [50–55]. As the hydrogen atoms exhibit  $\pm\frac{1}{2}$  spin, and the nuclear spins exist in two states that are randomly oriented, in absence of a net magnetic field, there is no overall net magnetization. When placed in an external magnetic field, the spins orient parallel and anti-parallel to the direction of the  $B_0$  magnetic field, with a slight propensity for the spins to align in the parallel direction, causing the tissue to express a net equilibrium magnetization [50, 53]. The magnetic moment of the atom rotates like a spinning top, predominately in the direction of the applied magnetic field. This magnetic moment rotates at an angular frequency unique to individual atoms, termed the Larmor frequency.

When a perpendicular radiofrequency field ( $B_1$ ) is applied at the hydrogen Larmor resonance frequency, only the protons absorb energy, and are tipped from the direction of the main magnetic field, with the flip angle denoting the degree that the spins are displaced from the equilibrium  $B_0$  direction [55]. This excites the protons to precess in a rotational motion around the  $B_0$  field vector. The excited proton magnetization vector then relaxes in the direction of the main  $B_0$  magnetic field, generating a longitudinal and transverse time-varying magnetization signal that is detected by the MRI receiver coils.

The rate at which this magnetization vector relaxes towards the main magnetic field direction is measured in terms of spin-lattice relaxation ( $T_1$ ) in the direction of the  $B_0$  magnetic field, and the spin-spin relaxation rate ( $T_2$ ) transverse to the  $B_0$  magnetic field direction [50, 53]. The  $T_1$  and  $T_2$  decay rates result from random static magnetic field variations. However, the relaxation rates are also influenced by time varying factors, such as magnetic field inhomogeneities, that combine with tissue static magnetic field to affect the relaxation rate.

The net magnetism applied to each proton results from both the field generated from the MRI system, in addition to the fields generated by the surrounding protons and bulk susceptibility [55, 56]. A chemical shift in the precession frequency results from the magnetic fields generated from these surrounding protons. This can allow identification of specific molecules present in the tissue, that introduce a distinctive chemical shift in the MR signal [55]. The degree of this shift also has a temperature dependence. Using the principle of proton resonance frequency shift (PRFS) thermometry, the individual temperature of each voxel can be quantified from the resulting temperature-dependent phase change due to this chemical shift [56].

## **4. MRI liver imaging**

Radiological imaging is used in a variety of manners in treating CRLM: including, to diagnose a condition, stage the disease, to locate extra-hepatic metastases, for treatment planning, for interventional image-guided procedures, and for post-treatment evaluation [57]. MRgHIFU requires additional MRI sequence protocols, compared to general diagnostic MRI.

### **4.1 Diagnostic MRI for CRLM**

Although CRLM is usually confirmed with computed tomography, MRI is an acceptable and common alternative, and is advantageous at identifying small lesions [1]. Some studies have shown MRI to provide the best results among all diagnostic imaging modalities, though more expensive [58]. The primary objectives for MRI liver tumor diagnosis are to verify the neoplasm presence, staging the lesion, and

classifying the type of neoplasm [22]. Accurate assessment of these techniques is crucial to guiding subsequent treatment such as resection, biopsy, and chemotherapy [22]. National Comprehensive Cancer Network (NCCN) guidelines recommend CT be used for initial workup and staging; with MRI recommended for potentially resectable cases, prior to locoregional treatment, and for inadequate imaging with CT [59].

Metastatic liver tumors have been reported as a factor of 18–40 more frequent than primary tumors [17]. The presence of both benign and malignant liver lesions are common. The challenge is often distinguishing the benign liver lesions from malignant lesions, as misdiagnosis can greatly impact staging and treatment planning. CRLM lesions exhibit T<sub>1</sub> signal hypointensity, higher FATSAT-T<sub>2</sub>W signal intensity, and higher diffusion-weighted imaging (DWI) signal intensity. On T<sub>2</sub>W, the tumor resembles a target; with coagulative necrosis causing a relatively higher signal intensity in the tumor center, followed by a reduced signal exterior due to bulk desmoplasia, and an even lower intensity thin edge from desmoplasia growing at the periphery. This thin edge resembles a ring in the arterial phase when gadolinium is administered. These features can change due to fatty liver infiltration and edema [60].

Standard liver tumor protocols are concerned with imaging the parenchyma, vascular supply, and biliary tract [61]. Basic liver protocols often include: T<sub>2</sub> half acquisition single-shot turbo spin echo (HASTE) localizer, in-op phase T<sub>1</sub> Gradient Recall Echo (GRE), T<sub>2</sub> fast spin echo (FSE) with fat saturation (FATSAT), and gadolinium-enhanced 3D FATSAT T<sub>1</sub> GRE [61, 62]. The HASTE localizer uses a motion insensitive T<sub>2</sub> single-shot spin echo sequence in combination with half-Fourier to acquire a multislice image in about 2 seconds during a single breath hold [63]. The in-op phase Dixon technique, is a spectroscopic technique used to suppress fat signal, quantify the hepatic fat content of the liver, and estimate iron content [63]. The spectroscopic method distinguishes an image at the  $-\text{CH}_2$  fat chemical shift from an image at the water chemical shift [64]. In-phase and op-phase sequences are often spin-echo or GRE sequences with equal repetition times, but different echo times. It acquires a normal in-phase image containing the water and fat, an opposed-phase image containing the water phase signal lessened by the fat phase contribution. Combining in-phase and op-phase images generates the water only image, and subtraction of the op-phase image from the in-phase image allows isolation of the fat signal [64, 65]. Additionally, the Dixon technique allows the generation of a T<sub>2</sub><sup>\*</sup> map, from which the local iron content (mg g<sup>-1</sup>) can be formulated [65, 66].

Of high importance in clinical diagnosis of liver lesions are DWI and hepatocyte-specific magnetic resonance contrast agent imaging, with MRI elastography to a lesser extent [67]. DWI is particularly useful for detection of small metastatic lesions [61]. Liver DWI consists of a T<sub>2</sub> sequence with symmetric diffusion sensitizing gradients centered on the 180° refocusing pulse [67, 68]. Brownian motion of water molecules is more restricted in tumors and provides a noticeable degree of contrast compared to normal tissue [69]. The DWI sequence is generally used without the administration of a contrast agent, making it a completely non-invasive diagnostic sequence. The weighting factor in DWI is adjusted based on the b-value, that is a function of gradient strength and duration. The apparent diffusion coefficient (ADC) maps can be viewed by removing the T<sub>2</sub>-weighting from a series of diffusion-weighted images. Hyperintense regions generally correspond to regions of low fluid diffusion [63].


CRLM lesions are a solid liver lesion and a general protocol for identification and characterization can be described as follows. First, a highly T<sub>2</sub>-weighted SSTSE to identify benign fluid-filled lesions, such as cysts and hemangiomas. Next, a modestly T<sub>2</sub>-weighted FATSAT-TSE or DWI to identify metastatic tumor sites. Then, a Dixon

sequence might be used to observe the degree of fat infiltration into the tumor. Lastly, a contrast-enhanced image can be used for  $T_1$ -weighted phase imaging to characterize the tumor [70].

Extracellular gadolinium agents are the most common contrast agents for general imaging throughout the body [71]. Two common hepatic specific contrast agents are gadoxetate disodium (Gd-EOB-DTPA, Primovist, Eovist, Bayer Healthcare Pharmaceuticals) and gadobenate dimeglumine (Gd-BOPTA, Bracco Diagnostics) [67]. In some studies, Gd-EOB-DTPA hepatocyte specific MRI contrast agents has shown improved sensitivity and specificity in diagnosis of liver metastasis compared to computed tomography, particularly to the improved ability to detect small metastases [67, 72]. The hepatic-specific contrast agents are specific to tumors originating from hepatocytes, and can help distinguish these lesions from cavernoma or metastatic lesions [71, 73]. Though, these are more expensive than extracellular analogues, have a lower recommended dose and signal, and can exhibit reduced uptake in patients with hepatocyte dysfunction [69]. A comparison of DWI and Gd-EOB-DTPA- $T_1$ W MRI for detecting small lesions from CRLM are shown in **Figure 1** [74].

#### 4.2 MRgHIFU sequence aspects

MRgHIFU requires additional MRI sequences that allow for temperature mapping. MR temperature mapping most commonly utilizes PRFS thermometry [75–77], though other possible techniques allow temperature measurements based on the temperature-dependence of relaxation rates, proton density, water diffusion coefficient, thermo-sensitive contrast agents, and magnetization transfer [78, 79]. Resonance frequency shift results from temperature differences in water molecules and aqueous tissues, due to varying degrees of hydrogen bonding. At increased temperatures, the amount of hydrogen bonding is reduced. This increases nuclear shielding of water protons from the incident magnetic field, generating a lower resonance frequency in the water molecules [78]. This results in a linear-dependence of the phase map values from the chemical shift due to temperature change, at a rate of about  $-0.01$  ppm  $^{\circ}\text{C}^{-1}$  [78, 79]. MRgHIFU sequences



**Figure 1.** Comparison of diffusion-weighted MRI with contrast-enhanced  $T_1$ W MRI. Left: diffusion-weighted MRI of liver metastases. The arrows indicate small metastatic tumors, less than 1 cm diameter. Right: CE- $T_1$ W image after applying Gd-EOB-DTPA, in the same patient. Reprinted with permission from Koh and Berry [74].

are often based on GRE segmented echo-planar imaging (SEG EPI) sequences. A basic GRE sequence is the fast low-angle shot (FLASH) sequence that utilizes small flip angles to obtain a short echo time (TE) and repetition time (TR) [80]. The sequence further benefits from EPI to accelerate the acquisition rate.

Additional sequences are used to assess tissue peri-ablation and post-ablation. During the peri-ablation period, inflammation in the focal region results from edema, giving more contrast enhancement, and remains for some months. After ablation,  $T_2$  and peripheral  $T_1$  hyperintensity increases significantly due to the presence of hemorrhagic debris at the ablation region. Thickening or nodule formation in the peripheral hyperintense signal can also indicate recurrence or incomplete ablation, during the months following the procedure [67]. Alternative sequences are under study for other aspects of the modality. For example, magnetic resonance acoustic radiation force impulse (MR-ARFI) sequences allow simultaneous displacement and temperature measurements [81, 82], and is implemented in clinical research settings for tracking focal spot and assessing positioning errors [83, 84]. Additionally, MR-ARFI sequences are being studied for phase aberration correction that occurs in transcostal or transcranial procedures [85–87].

Also, thermal ablation needs temperature processing less than about one second. The faster sequences result in reduced signal to noise ratio and increased temperature uncertainty. Echo planar imaging, parallel imaging, alternate trajectories, and undersampling can increase the MRI frame rate [88–90]. In typical rectilinear sampling, the RF pulse frequency and slice-select gradient determine the slice to be imaged, the frequency encoding gradient amplitude controls the  $k_x$ -dimension position, and the phase encoding gradient amplitude controls the  $k_y$ -dimension position [54]. Alternate trajectories are useful, particularly for fast acquisition times and reducing motion artifacts. Radial trajectories are utilized in some of the fastest real-time MRI sequences [90]. Magnetic field inhomogeneities and magnetic susceptibility are also significant aspects to proper imaging and temperature mapping [91, 92].

## 5. Surgery for CRLM

Primary colon cancer is classified IV in patients presenting CRLM [2]. Most CRLM patients develop liver metastases after initial CRC treatment, while about 20–34% present liver nodules at initial diagnosis [59]. When CRLM are confined to the liver, the intent should be cure, and surgical resection is actually the standard of care, presenting the best survival rates [59, 93]. Neoadjuvant and adjuvant systemic therapy are recommended in most patients prior to surgical resection, as it can improve instances of recurrence [1, 59]. The response to neoadjuvant chemotherapy has shown to be a strong prognostic factor for outcomes after hepatic resection [94]. The aim of liver resection is to remove all macroscopic disease with clear (negative) margins and leave sufficient functioning liver, with proper vascular and biliary flow [95]. An inadequate future liver remnant volume (FLRV) can lead to post-hepatectomy liver failure, a major cause of morbidity and mortality. Typically, FLRV is intended to be more than 30% of the native tissue and 30% future liver remnant, or more than 350 grams of liver remaining per 70 kg body weight [1]. The anatomic description of functional segments, which is based on the organ's blood supply via the hepatic artery and portal vein, its venous drainage via the hepatic veins, and lastly its biliary drainage, is the foundation of liver surgery. Historically, up to six Couinaud segments can be removed in healthy individuals, returning to original size in about three weeks,




with restored liver function in about six weeks [29, 96]. An illustration of the liver segments are given in **Figure 2**.

Typical resection complications occur in 20–50% of patients, although the mortality rate is only 1–3% in high volume centers [29, 97]. Most common complications include pleural effusion or pulmonary atelectasis, venous catheter infection, site-incisional infection, ascites, subphrenic infection, intraperitoneal bleeding, biliary tract hemorrhage, coagulation disorders, and bile leakage [98]. Additionally, inadequate post-operative liver response can result from pre-operative liver dysfunction, prolonged vascular occlusion, and inadequate residual liver volume; leading to hepatic insufficiency that results in ascites, mental impairment, hyperbilirubinaemia, and possible sepsis [98]. Post-operative liver function can be evaluated by dynamic functional testing such as indocyanine green (ICG) clearance rate, or by aminopyrine breath tests for cytochrome P-450 function, and post-treatment monitoring with blood serum tests for analytes including coagulation products and albumin [98].

Surgical resection for synchronous CRLM is an extremely complex scenario and surgery remains one of the major curative treatment options available.

Consideration for surgical resection must be given to: the anatomical distribution of the disease; FRLV; management of the primary disease (in the setting of synchronous CRLM); the timing and role of (neo)adjuvant chemotherapy, and whether all disease can be resected successfully at one sitting. Patients are often administered chemotherapy and chosen to undergo a conventional colon-first procedure, a liver-first procedure, or simultaneous resection [99]. Even for patients presenting multifocal bilateral CRLM, the goal should be a full tumor excision with sufficient remaining functional parenchyma. Though, for multifocal bilateral CRLM, resection and ablation often yield survival rates only faintly superior to chemotherapy alone [99]. The traditional colon-first approach involves complete primary CRC tumor resection, along with systemic chemotherapy, then hepatectomy is performed later if resectable [100]. A “liver-first” approach involves




**Figure 2.** Illustration identifying locations of individual Couinaud segments. Olga Bolbot/shutterstock.com.

initial systemic chemotherapy, liver tumor removal, then CRC resection [100]. The concept is that the liver tumor is most likely to create further metastasis and the CRC is quite sensitive to systemic chemotherapy [101]. With either approach, approximately only 10–20% of patients are surgical candidates [2, 8, 102]. Reasons include late-stage cancer diagnosis, secondary tumor sites outside the liver, and existing comorbidity ineligibility [2, 8]. Although surgical resections report long-term survival rates, about half of the patients develop widespread metastases within three years [1]. Recurrence after primary liver resection occurs at about a 43% rate in the liver and about a 31% rate in the lungs [8].

Anatomic resections usually involve two or more hepatic segments, while non-anatomic resection involves resection of the metastases with a margin of uninvolved tissue (segmentectomy). Various approaches in liver resection include: right hepatectomy, right lobectomy, left hepatectomy left lobectomy, extended right hepatectomy, and extended left hepatectomy [103]. By performing a segment-based resection, intraoperative hemorrhage and remaining post-treatment ischemic tissue can be avoided, helping to prevent infection and bile duct fistula. Additionally, the segment-based approach allows predetermined calculation of tumor margins and remaining viable parenchyma. Moreover, intrahepatic metastases tend to arise in the same Couinaud segments, allowing better chances to remove small satellite metastatic sites [103, 104].

Modern surgery resection is based on the report of the first successful procedure for a right hepatectomy [103, 105]. An illustration of the basic liver anatomy is shown in **Figure 3**. Each Couinaud segment is functionally independent, receiving blood supply from the portal vein and from the hepatic artery; at the same time the outflows is guaranteed by various branches of the hepatic vein. The right hepatic lobe is composed of Couinaud segments 5–8, with the blood supply to the right lobe provided by the right portal vein and right hepatic artery. First, the falciform ligament, coronary ligament, and right triangular ligament are cut to allow increased liver movement. Next, the right hepatic artery, right portal vein, right hepatic duct, and cystic duct are clamped, cut, and ligated. The blood supply to the left lobe is kept intact. This is followed by dissection of the right lobe from the inferior vena cava. Venous outflow from the main and short hepatic veins are divided and ligated. This devascularization creates a line of demarcation due to a color change in the right liver lobe. Then,



**Figure 3.**  
*Overview of the liver anatomy. Olga Bolbot/shutterstock.com.*

transection of the liver parenchyma occurs, dividing the right and left lobes along the middle hepatic vein. This is followed by ligation of the middle hepatic vein blood supply. Parenchyma transection can result in large blood loss that can be lessened using a reduced central venous pressure and Pringle's manoeuvre [103, 104, 106]. Three major complications for the procedure include the introduction of an air embolism into the hepatic veins, hemorrhagic bleeding from the hepatic veins, and biliary leakage into the abdominal cavity [107].

The concept of the "two-staged hepatectomy" has been introduced by Adam et al. [108], as a surgical strategy that could be applied to patients with conventionally irresectable metastases to make them eligible for liver resection. This approach involved a combination of systemic chemotherapy to downstage tumors, with or without portal vein embolization (PVE), with subsequent planned staged operations that permitted curative resection of large tumor burden that would otherwise have been considered unresectable. The interval between operations enabled hypertrophy of the remnant liver to theoretically reduce the chance of liver insufficiency and patients would receive chemotherapy during the interval between operations in an effort to control tumor growth.

More recently, the technique known as ALPPS (Associating Liver Partition and Portal Vein Ligation for Staged Hepatectomy) allows removal of extensive tumor load by increasing future liver remnant, allowing increased surgical eligibility, and extended survivability of CRLM patients [109, 110]. Early research included right PVE, which was shown to induce hypertrophy in the left lobe, subsequently allowing increased amounts of liver tissue to be removed in the right lobe [108, 110–112]. This was later applied in two-stage hepatectomies to allow increased amounts of cancerous liver tissue to be removed from both liver lobes, by permitting liver regrowth between procedures [110]. Early two-stage hepatectomies required months for liver regrowth, with tumor progression frequently occurring during this time; however, development of ALPPS allowed the two surgical procedures to be performed within 7–14 days [109, 110]. ALPPS is indicated in case of extensive multifocal CRLM, failure after portal vein embolization, and expected small amounts of FLRV [14].

A generic procedure for two-stage hepatectomy of left lobe wedge resection combined with right lobe hepatectomy includes *in situ* liver splitting in addition to portal vein ligation [113]. First, the falciform ligament is cut, then tumors locations are confirmed and marked by intraoperative ultrasound. The transection line(s) is identified. Then, the right cystic duct and artery are ligated, followed by dissection and ligation of the right portal vein at the portal bifurcation. The right and middle hepatic veins are isolated, the space between is dissected, and umbilical tape is placed for the hanging maneuver. Then, transection of the parenchyma is performed at the site previously marked with/without Pringle maneuver. The liver is patched, drains placed, abdomen closed, ending the first stage. At this stage the liver is separated but not removed. Then, functional liver testing and weekly volumetry measurements are performed with CT or MRI until the future liver remnant volume surpasses 30%. In the second stage, the incision is reopened, and the hepatic artery and bile duct are ligated on the right lobe that previously underwent portal vein ligation. Then, transection of the right hepatic vein is followed by removal of the right liver lobe and closure of the abdomen.


In the last decade, it has been conceptualized that liver transplantation could offer the theoretical advantage of a real R0 resection, removing also all potentially undetected metastases. Earlier studies in American and European populations showed that transplant after non-neuroendocrine liver metastases from various primary

sites yielded one-year survival rates of only 5%, which is compounded by the lack of available donors [33]. More recent studies with tightened inclusion criteria have shown more favorable outcomes and resulted in a large increase in CRLM transplants worldwide [30, 31, 114]. The studies have suggested much longer survival rates after liver transplant for CRLM, when the inclusion criteria included adequate response to chemotherapy, excised primary tumor sites, more than one year between diagnosis and transplant, and liver only metastases [31, 32, 115]. Additional exclusion criteria exist based upon molecular profiling; for instance, exclusion is recommended due to V600 BRAF mutations and MSI from DNA mismatch repair (MMR) mutations [116]. These results have suggested liver transplant possibly provides the best overall survivability compared to other treatment modalities for surgical ineligibility. The drawbacks are smaller study size, the limited availability of liver donors and more specialized training is required across multiple disciplines to conduct the operation [30].

## 6. Chemotherapy for CRLM

Systemic chemotherapy in CRLM is administered to attain surgical eligibility, for disease control, peri-operatively, or palliatively; since the treatment alone is rarely curative, with 5-year survival rates less than 10%, and historically less than 1% [25, 30, 117]. Polymetastatic liver disease faces treatment limitations with chemotherapy being the primary treatment. The survival rate is poor and a large demand exists for improved treatment options. As surgical resection offers the best long-term survival rates, the aim of systemic chemotherapy is often to downsize tumors to convert ineligible patients to surgical candidates, with systematic review showing a conversion rate for R0 resection in initially ineligible patients at 23% [118]. Chemotherapy regimens are administered neoadjuvantly prior to hepatectomy for cytoreduction, to reduce metastatic tumor size, allowing smaller resection volumes [119]. The regimens are also administered after resection to reduce recurrence [25, 120]. Hepatic intra-arterial infusion is often beneficial because the liver metastasis is supplied by the hepatic artery network, normal tissue is supplied by the portal vein, and locoregional treatment can be performed without exposing much healthy tissue [121–123]. The liver contains a capillary network of sinusoids that filter the blood as shown in **Figure 4**. Approximately 45% of metastatic tumor cells, predominately arriving from the hepatic arterial network [123], become embedded in the sinusoids [89]. Normal liver parenchyma receive about 80% of the blood supply from the portal vein and about 20% from the hepatic artery. In contrast, about 80% of the tumor blood supply arrives from the hepatic artery [116]. This allows locoregional embolization techniques, like radioembolization and chemoembolization, to both embolize the blood supply to specific tumor segments, and deliver locoregional radiotherapy or chemotherapy. These embolization techniques are suggested to be considered for metastatic CRC limited only to the liver, and after unsuccessful chemotherapy [1].

The chemotherapy regimen depends on a number of factors, including: aim of cytoreduction prior to surgery, aim of disease control, aim of palliation, type of somatic gene mutation, and wild-type or mutant phenotype. Somatic mutations of RAS proto-oncogenes have been found in up to 52% of CRLM hepatic resections, with up to 6–12% of resections expressing BRAF mutations, and co-occurring proto-oncogene RAS and TP53 tumor-suppressor mutations as common genetic events [1, 94]. According to ESMO guidelines, first-line chemotherapy for cytoreduction in RAS



**Figure 4.** *Histological depiction of liver lobules. These units are microscale components of liver tissue. The liver sinusoids are small capillaries, with blood supplied by small branches of the hepatic artery and portal vein. Design/shutterstock.com.*

tumors should be recommended cytotoxic doublets (FOLFOX/CAPOX/FOLFIRI), in combination with VEGF antibody bevacizumab for RAS mutant-type tumors, and EGFR antibodies for wild-type tumors. FOLFOXIRI with bevacizumab are recommended as a first line treatment for cytoreduction in CRLM BRAF mutant tumors [1].

Chemotherapeutics can also exhibit many adverse side-effects on healthy liver tissue. Side-effects include sinusoidal obstruction syndrome and chemotherapy-associated steatohepatitis, that can lead to liver failure or increased mortality rates [119]. Additionally, the chemotherapy can cause missing metastases, making lesions unidentifiable on radiological imaging, complicating surgical decisions, and increasing the chance of recurrence [119]. Chemotherapy has difficulty supplying tumor cells with adequate drug dose. The maximum dose is limited by systemic toxicity effects and inadequate tumor penetration is common [8, 124]. Intrahepatic arterial delivery can exhibit acute side-effects of hepatocellular atrophy causing cirrhosis and necrosis [123, 125].

Doxorubicin is an anthracycline chemotherapeutic that can be administered during combination therapy. A liposomal form was created relatively early due to the need for better treatment in Kaposi sarcoma from autoimmune deficiency syndrome [126]. Clinical trials of FUS-mediated thermosensitive liposomal doxorubicin drug delivery to liver tumors [127, 128] have shown large increases in intratumoral doxorubicin concentration, and there are ongoing trials with MRgHIFU for pediatric tumors [129]. Similar ongoing trials are studying the enhanced ability for microbubbles to improve chemotherapy delivery to metastatic liver tumors [130].

## 7. Radiotherapy for CRLM

Radiotherapy emits ionizing radiation at tumors, causing DNA damage, and apoptosis. The technique exhibits some similar drawbacks to focused ultrasound. Cumulative radiation exposure can occur in the beam's near and far field, resulting in unwanted tissue damage [8, 131, 132]. Also, systems require computed tomography

guidance and respiratory motion control [8, 133]. Local ablative techniques, including radiotherapy, are generally considered to be limited to patients with unresectable CRLM or oligometastatic disease [1]. CRLM radiotherapy has often been limited by liver parenchyma radio-sensitivity. External beam radiation doses of 70–90 Gy needed for CRLM and HCC tumor treatment exceeds tolerance limits of 35 Gy for radiation-induced liver disease (RILD) [57, 134] that can lead to liver failure and death [25]. The condition occurs two to sixteen weeks after treatment, is identified by ascites, high levels of alkaline phosphatases, and high levels of liver transaminases [135].

Stereotactic body radiation therapy (SBRT) with linear accelerators has recently gained much interest for surgical ineligibility, particularly in oligometastatic disease. With SBRT, fiducial markers are percutaneously placed near the tumor site to allow precise tumor targeting [57]. Though MRI guidance reduces invasiveness, without the need for fiducial markers [116]. SBRT is recommended by ESMO to be considered for patients with oligometastatic disease who are ineligible for surgery and ablative therapy [1]. One major advantage of SBRT compared to ablative therapies is that the treatment is non-thermal, mitigating some of the common side-effects seen in local ablative techniques, such as fluid perfusion effects [116]. Studies have shown that liver failure is infrequent when only a portion of the liver is irradiated [135]. The liver toxicity is mild to moderate, with liver failure in less than 1% of patients [136, 137]. Treatment of oligometastatic CRC in the liver with SBRT, suggests one and two year overall survivability at about 67.1% and 56.5%, respectively [137]. Many early phase clinical trials are recruiting, active, or recently completed, for treatment of primary or secondary hepatic tumors with magnetic resonance guided linear accelerators [138, 139] and magnetic resonance guided SBRT [140–143]. Recent phase I trial results with magnetic resonance guided SBRT, showed improved toxicity, with estimated 2-year overall survival of 51%, and median overall survival of 29 months [144].

## **8. Focused ultrasound clinical studies for liver cancer**

A substantial number of clinical studies, cohorts, and randomized control trials for non-liver MRgHIFU and MRgFUS have been reported, including: treatment with bone osteomas or palliative bone metastasis [145, 146], uterine fibroids [147–151], gynaecological tumor recurrence [152], prostate cancer [153, 154], essential tremor [155, 156], and breast cancer [157]. Many clinical studies have been reported for USgFUS ablation for liver tumors [158–165], with most studies reporting on HCC ablation [166]. Similar to USgFUS, new histotripsy devices using cavitation rather than thermal ablation, are currently being studied for the treatment of primary and secondary tumors, with an active prospective clinical trial [45, 46, 167, 168]. No Phase III trials for USgFUS or MRgHIFU ablation of CRLM have been published [116]. Early USgFUS studies in liver malignancies, not distinguishing between metastatic liver tumors and primary liver tumors, showed a median survival time of 13.4 months, 6-month survival times of 82.6%, and 12-month survival time of 53.4% [159]. More recent systematic reviews of FUS for liver malignancies have given 1 year, 2 year, and 5-year survivability of 81%, 60%, and 39%, respectively [166]. Most studies have been conducted using the Chongqing Haifu JC system, capable of up to 300 W acoustic power and peak intensity up to 20,000 W cm<sup>2</sup> [166]. The system has received the mark *Conformite' Europeenne' (CE)*, being the most reported system for clinical liver tumor ablation [2, 3]. The permission is granted to individual commercial models rather than general treatment procedures. The magnetic resonance guided

systems that have received regulatory approval for alternative treatments include the ArcBlate (Episonica, Hsinchu, Taiwan), Exablate (Insightec, Tirat Carmel, Israel), and Sonalleve (Profound Medical, Mississauga, Canada) systems.

Local ablative techniques, including focused ultrasound ablation, are generally recommended only in cases of unresectable liver metastases or oligometastatic disease [1]. Most FUS ablation therapy studies for liver tumors are USgFUS for HCC, with less reports of metastatic liver tumor treatment [166]. Particularly advantageous in FUS is the improved side effect profile and reduced morbidity compared to standard treatment options. The treatment can occur multiple times with no cumulative radiation-like side effects. In relation to chemotherapy, it is much more focused, with less toxicity to healthy tissues [8, 169]. Additionally, extracorporeal FUS liver ablation is completely non-invasive and offers very fast recovery times [170]. Benefits of MRgHIFU compared to USgFUS include near real-time temperature mapping, integration into existing imaging systems, less propensity for radiofrequency interference in the imaging system, and capability of assessing treatment response during the procedure. Though ultrasound-guided devices do not provide real-time temperature mapping, assessment of grey-scale change are indicative of coagulative necrosis [166]. Treatment plans with FUS generally depend on the cancer staging. Curative ablation of early stage tumors often include a 1.5–2.0 cm peripheral tissue margin. The treatment is administered palliatively for late-stage tumors to slow progression or alleviate symptoms [8, 160].

Drawbacks to hepatobiliary focused ultrasound studies have been the need for general anesthesia, long treatment times, scattering by the thoracic cage, high power requirements, respiratory motion, skin burns, osteonecrosis, skin pain, skin edema, rib resection, fever, the need for intrapleural effusion, and reduced thermal dose from fluid perfusion of surrounding vessels [2, 39, 165, 166, 170–177]. A systematic review of USgFUS for the treatment of malignant hepatobiliary tumors indicated the primary complications were skin burns in 15% of cases, followed by localized pain in 5%, then fever at 2% [166]. Major post-treatment complications include fluid and/or air accumulation in the lungs, biliary obstruction, and fistula occurrence [177].

Some studies have reported focused ultrasound ablation in primary and secondary liver tumors in difficult locations, including near major hepatic veins and arteries, and near surrounding organs of the heart, gallbladder, stomach, and intestine [162, 165, 178]. Tumors located near surrounding organs are high-risk. Particularly sensitive are the bowel and gallbladder due to the thin walls and risks of peritonitis [162].

Skin and rib burns have been addressed in a variety of manners. Skin burns have been reported to occur with tumors located near the subcapsular area, resulting from possible rib reflection or reflections from internal gas pockets in the bowel or lung parenchyma [166]. The right lobe is more susceptible as it is predominately located behind the ribs [162]. Intrapleural effusion can distance the tumor site from the subcapsular area, or rib resection can be performed [162, 179]. Particularly troublesome are tumors of the liver dome in Couinaud segments 7 and 8, due to the close proximity to the lungs, the close proximity to the ribs, and that this region tends to remain behind the rib cage under general anesthesia due to reduced respiration [162]. A small cohort for USgFUS reported that proper intraoperative assessment of the soft tissue prevented skin burns in all patients [161].

A variety of techniques have been tested to overcome respiratory motion and rib interaction. Respiratory motion creates complications requiring organ image

registration techniques [180] and MRI motion artifact compensation [174, 181]. Numerous preclinical studies have undertaken new technologies to address respiratory motion and rib interactions [39, 172, 174–176, 180, 182–186]. Previous USgFUS human studies have generally been successful at performing ablation through the ribs; though additional measures have included left lung ventilation with endotracheal intubation and general anesthesia to reduce liver movement, intrapleural effusion, and rib resection [160, 162, 179]. MRgHIFU pilot studies used intermittent sonications, and limited to the treatment to the left liver lobe, in tumor sites not blocked by the ribs [170, 187–189].

Handheld intraoperative HIFU devices under ultrasound-guidance are in development, and being tested in early phase clinical trials for CRLM tumor ablation. The technique is similar to intraoperative radiofrequency and microwave ablation, but prevents the need for an intraparenchymal probe. Results have been reported using the device for ablating tissue near tumors in segments prior to surgical resection, to assess accuracy and safety. Applications include reduction of hemorrhaging during surgery and potentially bridging more patients for surgical resection [190–193]. The device was shown capable of in vivo hepatic vessel occlusion for diameters of 2 mm [194], and studies have reported diameters of left hepatic arteries and right hepatic veins between 3 and 4 mm [195].

Several small clinical studies have been reported for MRgHIFU ablation for HCC [170, 187–189, 196, 197]. There is currently an ongoing Phase I clinical trial with MRgHIFU for a variety of pediatric solid tumors, in which hepatic tumors are eligible [198].

In the study from Okada et al. [187], MRgHIFU liver tumor ablation was performed on a single patient. The MRI system utilized respiratory gating and ablation was performed on a 15 mm HCC lesion. The procedure required about two hours to ensure complete coagulation by repeated coverage. Gadolinium contrast agent was administered post-treatment and no increased signal intensity was observed at the tumor site, indicating expected ablation contrast. The authors noted the need for better technology for avoiding bowel loops, ribs, and respiratory liver motion. Though, the patient only complained of slight skin heating discomfort during treatment and was released from the hospital the following day.

Anzidei et al. treated a single HCC patient more comprehensively with MR-FUS [188]. The patient refused surgery and percutaneous ablation, then opted for MR-FUS. The individual had no distant metastases, was treated successfully, and later underwent total liver transplant. Excised liver histopathology showed complete coagulative necrosis with only slight recurrence at the ablation periphery. The investigators noted the procedure can be improved with better respiratory motion control and expected that future applications would use automated feedback algorithms.

Gedroyc conducted a series of pilot studies for MRgHIFU liver tumor ablation [170, 189]. It was reported that the absorption from the ribs was problematic and the treatments were limited to patients with exposed tumor sites, such as below the rib line or in the left lobe of the liver. One case was a female with HCC arising from Hepatitis-B infection. She was previously treated with hepatic arterial chemoembolization and laser ablation. Recurrence occurred with a 1.5 cm HCC lesion in the left lobe within Couinaud segment 3, a position that was not covered by the ribs. The site was ablated with MRgHIFU. In another case, the patient was a male with HCC, Hepatitis-C, extensive cirrhosis, and elevated alpha-fetoprotein levels. He was treated for a 3 cm HCC in the anterior portion of the left liver lobe.



## **9. Conclusions**

Optimal treatment strategies for CRLM patients should be made by a multi-disciplinary team as part of a tumor board, for establishing diagnostic and treatment strategies [1]. Surgery of CRLM will likely provide the best long-term outcomes and the strategy should focus on complete resection. Although, the majority of patients with CRLM are ineligible for surgery and many surgical cases will experience widespread recurrence. Thermal ablation methods like focused ultrasound are generally only recommended for unresectable CRLM and oligometastatic disease, with at least one system under ultrasound guidance having the CE-mark for CRLM [1, 2]. Due to expected increasing CRLM incidence and high surgical ineligibility, non-invasive technologies like MRgHIFU systems have great potential for clinical translation as an ablative interventional radiology procedure.

Guidelines for FUS pilot studies suggest performing MRgHIFU ablation in CRLM patients prior to the surgical operation, then surgically removing the ablated tumor, and assessing the effectiveness with pathology [2]. Randomized control trials have been suggested to be performed on CRLM patients that are not candidates for surgical resection or RF ablation, and to compare TACE and MRgHIFU to a control group receiving only TACE [2]. In a randomized controlled trial, USgFUS for primary liver tumors in combination with TACE has shown improved treatment over TACE alone, increasing survival times, giving higher remission rates, lower recurrence rates, lower rates of post-operative metastases, and less instances of hemorrhaging in the digestive tract [199, 200].

MRgHIFU has been established in proof-of-concept studies for HCC, limited to the left liver lobe or section not covered by the ribs, requiring intermittent ablation due to respiration, and has not been reported in randomized controlled trials for primary or secondary liver tumors [170, 187–189, 196, 197, 201]. The FUS field has gained much interest in recent years and MRgHIFU ablation of primary and secondary liver tumors appears likely to begin early phase trials in the near future. Previous focused ultrasound studies have developed methods to address many technical complications such as respiratory motion and suppressing prefocal interactions; with focusing through the ribs being one of the major technical difficulties. Long treatment times are another complication and should improve with automated feedback control and faster acquisition times.

IntechOpen

## Author details

Ryan Holman<sup>1\*</sup>, Orane Lorton<sup>1</sup>, Pauline C. Guillemin<sup>1</sup>, Andrea Peloso<sup>2,3</sup>,  
Alexis Ricoeur<sup>4</sup> and Rares Salomir<sup>1,4</sup>

1 Image Guided Interventions Laboratory, Department of Radiology, University of Geneva, Geneva, Switzerland

2 Hepatology and Transplantation Laboratory, Faculty of Medicine, Department of Surgery, Division of Visceral Surgery, University of Geneva, Geneva, Switzerland


3 Department of Surgery, Division of Visceral Surgery, Geneva University Hospitals, Geneva, Switzerland

4 Radiology Department, University Hospitals of Geneva, Geneva, Switzerland

\*Address all correspondence to: [ryan.holman@unige.ch](mailto:ryan.holman@unige.ch)

## IntechOpen

---

© 2022 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. 

## References

- [1] Van Cutsem E, Cervantes A, Adam R, Sobrero A, Van Krieken JH, Aderka D, et al. ESMO consensus guidelines for the management of patients with metastatic colorectal cancer. *Annals of Oncology*. 2016;27(8):1386-1422
- [2] Aubry JF, Pauly KB, Moonen C, Haar G, Ries M, Salomir R, et al. The road to clinical use of high-intensity focused ultrasound for liver cancer: Technical and clinical consensus. *Journal of Therapeutic Ultrasound*. 2013;1(13):1-13
- [3] State of the Field Report. Focused Ultrasound Foundation. 2021. Available from: <https://www.fusfoundation.org/the-foundation/foundation-reports#SOTF>
- [4] World Health Organization (WHO). Global Health Estimates 2020: Deaths by Cause, Age, Sex, by Country and by Region, 2000-2019. WHO; 2020. Available from: <https://www.who.int/data/gho/data/themes/mortality-and-global-health-estimates/ghe-leading-causes-of-death>; [Accessed: December 31, 2021]
- [5] Cassidy J, Bissett D, Obe RAS, Payne M, Morris-Stiff G. *Oxford Handbook of Oncology*. OUP Oxford; 2015
- [6] Sung H, Ferlay J, Siegel RL, Laversanne M, Soerjomataram I, Jemal A, et al. Global Cancer Statistics 2020: GLOBOCAN Estimates of Incidence and Mortality Worldwide for 36 Cancers in 185 Countries. *CA: A Cancer Journal for Clinicians*. 2021;71(3):209-249
- [7] Shlomai A, de Jong YP, Rice CM. Virus associated malignancies: The role of viral hepatitis in hepatocellular carcinoma. *Seminars in Cancer Biology*. 2014;26:78-88
- [8] Lyon PC. Targeted release from lyso-thermosensitive liposomal doxorubicin (ThermoDox®) using focused ultrasound in patients with liver tumours [doctoral dissertation]; 2016
- [9] Bruix J, Sherman M. Management of hepatocellular carcinoma. *Hepatology*. 2005;42(5):1208-1236
- [10] Bruix J, Sherman M, Llovet JM, Beaugrand M, Lencioni R, Burroughs AK, et al. Clinical management of hepatocellular carcinoma. Conclusions of the Barcelona-2000 EASL conference. European Association for the Study of the Liver. *Journal of Hepatology*. 2001;35(3):421-430
- [11] Fan ST. Hepatocellular carcinoma—Resection or transplant? *Nature Reviews Gastroenterology & Hepatology*. 2012;9(12):732-737
- [12] Vogel A, Cervantes A, Chau I, Daniele B, Llovet JM, Meyer T, et al. Hepatocellular carcinoma: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Annals of Oncology*. 2018; 29:iv238-iv255
- [13] Clavien PA, Lesurtel M, Bossuyt PM, Gores GJ, Langer B, Perrier A. Recommendations for liver transplantation for hepatocellular carcinoma: An international consensus conference report. *The Lancet Oncology*. 2012;13(1):e11-e22
- [14] Petrowsky H, Fritsch R, Guckenberger M, De Oliveira ML, Dutkowski P, Clavien PA. Modern therapeutic approaches for the treatment of malignant liver tumours. *Nature Reviews. Gastroenterology & Hepatology*. 2020;17(12):755-772

- [15] Tsilimigras DI, Brodt P, Clavien PA, Muschel RJ, D'Angelica MI, Endo I, et al. Liver metastases. *Nature Reviews Disease Primers*. 2021;7(1):27
- [16] Abdalla EK, Adam R, Bilchik AJ, Jaeck D, Vauthey JN, Mahvi D. Improving resectability of hepatic colorectal metastases: Expert consensus statement. *Annals of Surgical Oncology*. 2006;13(10):1271-1280
- [17] Hussain SM, Sorrell MF. Colorectal metastases II: MRI findings in a fatty liver. In: *Liver MRI Correlation with Other Imaging Modalities and Histopathology*. 2nd ed. Berlin, Heidelberg: Springer; 2014. pp. 92-93
- [18] Fong Y, Blumgart LH, Cohen AM. Surgical treatment of colorectal metastases to the liver. *CA: A Cancer Journal for Clinicians*. 1995;45(1):50-62
- [19] Padman S, Padbury R, Beeke C, Karapetis CS, Bishnoi S, Townsend AR, et al. Liver only metastatic disease in patients with metastatic colorectal cancer: Impact of surgery and chemotherapy. *Acta Oncologica*. 2013;52(8):1699-1706
- [20] Ruiterkamp J, Ernst MF, de Munck L, van der Heiden-van der Loo M, Bastiaannet E, van de Poll-Franse LV, et al. Improved survival of patients with primary distant metastatic breast cancer in the period of 1995-2008. A nationwide population-based study in the Netherlands. *Breast Cancer Research and Treatment*. 2011;128(2):495-503
- [21] Adeyiga AO, Lee EY, Eisenberg RL. Focal hepatic masses in pediatric patients. *American Journal of Roentgenology*. 2012;199(4):W422-W440
- [22] Shelmerdine SC, Roebuck DJ, Towbin AJ, McHugh K. MRI of paediatric liver tumours: How we review and report. *Cancer Imaging*. 2016;16(1):21
- [23] Shimada H, Tanaka K, Endou I, Ichikawa Y. Treatment for colorectal liver metastases: A review. *Langenbeck's Archives of Surgery*. 2009;394(6):973-983
- [24] Faivre J, Manfredi S, Bouvier AM. Epidemiology of colorectal cancer liver metastases. *Bulletin de l'Académie Nationale de Médecine*. 2003;187(5):815-822; discussion 822-3
- [25] Mitchell D, Puckett Y, Nguyen QN. Literature review of current management of colorectal liver metastasis. *Cureus*. 2019;11(1):e3940
- [26] Van de Velde CJH. Treatment of liver metastases of colorectal cancer. *Annals of Oncology*. 2005;16:ii144-ii149
- [27] Bismuth H, Adam R, Levi F, Farabos C, Waechter F, Castaing D, et al. Resection of nonresectable liver metastases from colorectal cancer after neoadjuvant chemotherapy. *Annals of Surgery*. 1996;224(4):509-520; discussion 520-2
- [28] Creasy JM, Sadot E, Koerkamp BG, Chou JF, Gonen M, Kemeny NE, et al. Actual 10-year survival after hepatic resection of colorectal liver metastases: What factors preclude cure? *Surgery*. 2018;163(6):1238-1244
- [29] Braccia DP, Heffernan N. Surgical and ablative modalities for the treatment of colorectal cancer metastatic to the liver. *Clinical Journal of Oncology Nursing*. 2003;7(2):178-184
- [30] Glinka J, Ardiles V, Pekolj J, Mattera J, Sanchez Claria R, de Santibanes E, et al. Liver transplantation for non-resectable colorectal liver metastasis: Where we are and where we are going.

Langenbeck's Archives of Surgery. 2020;**405**(3):255-264

[31] Hagness M, Foss A, Line PD, Scholz T, Jørgensen PF, Fosby B, et al. Liver transplantation for nonresectable liver metastases from colorectal cancer. *Annals of Surgery*. 2013;**257**(5):800-806

[32] Dueland S, Grut H, Syversveen T, Hagness M, Line PD. Selection criteria related to long-term survival following liver transplantation for colorectal liver metastasis. *American Journal of Transplantation*. 2020;**20**(2):530-537

[33] Li K, Yang J, Liu C, Chen P. In: Yan L, editor. *Liver Resection of Secondary Liver Cancer*. Dordrecht: Springer Netherlands; 2016. pp. 153-157

[34] Hokland SL, Pedersen M, Salomir R, Quesson B, Stodkilde-Jørgensen H, Moonen CT. MRI-guided focused ultrasound: Methodology and applications. *IEEE Transactions on Medical Imaging*. 2006;**25**(6):723-731

[35] Lynn JG, Zwemer RL, Chick AJ, Miller AE. A new method for the generation and use of focused ultrasound in experimental biology. *The Journal of General Physiology*. 1942;**26**(2):179-193

[36] Fry WJ, Mosberg JWH, Barnard JW, Fry FJ. Production of focal destructive lesions in the central nervous system with ultrasound. *Journal of Neurosurgery*. 1954;**11**(5):471-478

[37] Lele PP. Concurrent detection of the production of ultrasonic lesions. *Medical & Biological Engineering*. 1966;**4**(5):451-456

[38] Tempany CMC, McDannold NJ, Hynynen K, Jolesz FA. Focused ultrasound surgery in oncology: Overview and principles. *Radiology*. 2011;**259**(1):39-56

[39] Lorton O, Guillemin PC, Mori N, Crowe LA, Boudabbous S, Terraz S, et al. Self-scanned HIFU ablation of moving tissue using real-time hybrid US-MR imaging. *IEEE Transactions on Biomedical Engineering*. 2018;**66**(8):2182-2191

[40] Mihcin S, Melzer A. Principles of focused ultrasound. *Minimally Invasive Therapy & Allied Technologies*. 2018;**27**(1):41-50

[41] Izadifar Z, Izadifar Z, Chapman D, Babyn P. An introduction to high intensity focused ultrasound: Systematic review on principles, devices, and clinical applications. *Journal of Clinical Medicine*. 2020;**9**(2):1-22

[42] Chavrier F, Chapelon JY, Gelet A, Cathignol D. Modeling of high-intensity focused ultrasound-induced lesions in the presence of cavitation bubbles. *The Journal of the Acoustical Society of America*. 2000;**108**(1):432-440

[43] Harper JD, Metzler I, Hall MK, Chen TT, Maxwell AD, Cunitz BW, et al. First in-human burst wave lithotripsy for kidney stone comminution: Initial two case studies. *Journal of Endourology*. 2021;**35**(4):506-511

[44] Pishchalnikov YA, Sapozhnikov OA, Bailey MR, Williams JJC, Cleveland RO, Colonus T, et al. Cavitation bubble cluster activity in the breakage of kidney stones by lithotripter shockwaves. *Journal of Endourology*. 2003;**17**(7):435-446

[45] Xu Z, Hall TL, Vlasisavljevich E Jr, FTL. Histotripsy: The first noninvasive, non-ionizing, non-thermal ablation technique based on ultrasound. *International Journal of Hyperthermia*. 2021;**38**(1):561-575

[46] Vidal-Jové J, Serres-Creixamés X, Ziemlewicz TJ, Cannata JM. Liver

- histotripsy mediated abscopal effect—  
Case report. *IEEE Transactions on Ultrasonics, Ferroelectrics, and Frequency Control*. 2021;**68**(9):3001-3005
- [47] Kennedy JE. High-intensity focused ultrasound in the treatment of solid tumours. *Nature Reviews. Cancer*. 2005;**5**(4):321-327
- [48] Prakash P, Diederich CJ. Considerations for theoretical modelling of thermal ablation with catheter-based ultrasonic sources: Implications for treatment planning, monitoring and control. *International Journal of Hyperthermia*. 2012;**28**(1):69-86
- [49] Sapareto SA, Dewey WC. Thermal dose determination in cancer therapy. *International Journal of Radiation Oncology, Biology, Physics*. 1984;**10**(6):787-800
- [50] Plewes DB, Kucharczyk W. Physics of MRI: A primer. *Journal of Magnetic Resonance Imaging*. 2012;**35**(5):1038-1054
- [51] Bloch F. Nuclear induction. *Physics Review*. 1946;**70**:460-474
- [52] Purcell EM, Torrey HC, Pound RV. Resonance absorption by nuclear magnetic moments in a solid. *Physics Review*. 1946;**69**:37-38
- [53] Radue EW, Weigel M, Wiest R, Urbach H. Introduction to magnetic resonance imaging for neurologists. *Continuum: Lifelong Learning in Neurology*. 2016;**22**(5):1379-1398
- [54] Westbrook C. *MRI at a Glance*. 3rd ed. Chichester, England: John Wiley & Sons; 2016
- [55] Storey P. In: Prasad PV, editor. *Introduction to Magnetic Resonance Imaging and Spectroscopy*. Totowa, NJ: Humana Press; 2006. pp. 3-57
- [56] de Senneville BD, Quesson B, Moonen CTW. Magnetic resonance temperature imaging. *International Journal of Hyperthermia*. 2005;**21**(6):515-531
- [57] Bajpai S, Kambadakone A, Guimaraes AR, Arellano RS, Gervais DA, Sahani D. Image-guided treatment in the hepatobiliary system: Role of imaging in treatment planning and posttreatment evaluation. *Radiographics*. 2015;**35**(5):1393-1418
- [58] Niekel MC, Bipat S, Stoker J. Diagnostic imaging of colorectal liver metastases with CT, MR imaging, FDG PET, and/or FDG PET/CT: A meta-analysis of prospective studies including patients who have not previously undergone treatment. *Radiology*. 2010;**257**(3):674-684
- [59] National Comprehensive Cancer Network. *NCCN Clinical Practice in Oncology: Colon Cancer*. NCCN.org 2021, version 3. Available from: <https://www.nccn.org/professionals/physiangls/pdf/colon.pdf>
- [60] Hussain SM, Sorrell MF. Colorectal metastases I: Typical lesions. In: *Liver MRI Correlation with Other Imaging Modalities and Histopathology*. 2nd ed. Berlin, Heidelberg: Springer; 2014. pp. 90-91
- [61] Donato H, Franca M, Candelaria I, Caseiro-Alves F. Liver MRI: From basic protocol to advanced techniques. *European Journal of Radiology*. 2017;**93**:30-39
- [62] Wile GE, Leyendecker JR. Magnetic resonance imaging of the liver: Sequence optimization and artifacts. *Magnetic Resonance Imaging Clinics of North America*. 2010;**18**(3):525-547, xi
- [63] Runge VM, Nitz WR, Heverhagen JT. *The Physics of Clinical MR Taught*

Through Images. 4th ed. New York: Thieme; 2018

[64] Dixon WT. Simple proton spectroscopic imaging. *Radiology*. 1984;**153**(1):189-194

[65] Hussain SM, Sorrell MF. Dixon-based sequence: Assessment of liver fat and iron deposition. In: *Liver MRI Correlation with Other Imaging Modalities and Histopathology*. 2nd ed. Berlin, Heidelberg: Springer; 2014. pp. 266-267

[66] Wood JC, Enriquez C, Ghugre N, Tyzka JM, Carson S, Nelson MD, et al. MRI R2 and R2\* mapping accurately estimates hepatic iron concentration in transfusion-dependent thalassemia and sickle cell disease patients. *Blood*. 2005;**106**(4):1460-1465

[67] Karaosmanoglu AD, Onur MR, Ozmen MN, Akata D, Karcaaltincaba M. Magnetic resonance imaging of liver metastasis. *Seminars in Ultrasound, CT, and MR*. 2016;**37**(6):533-548

[68] Liu PS. Liver mass evaluation in patients without cirrhosis: A technique-based method. *Radiologic Clinics of North America*. 2015;**53**(5):903-918

[69] O'Neill EK, Cogley JR, Miller FH. The ins and outs of liver imaging. *Clinics in Liver Disease*. 2015;**19**(1):99-121

[70] Hussain SM, Sorrell MF. Solid liver lesions. In: *Liver MRI Correlation with Other Imaging Modalities and Histopathology*. 2nd ed. Berlin, Heidelberg: Springer; 2014. p. 87

[71] Fidler J, Hough D. Hepatocyte-specific magnetic resonance imaging contrast agents. *Hepatology*. 2011;**53**(2):678-682

[72] Kim HJ, Lee SS, Byun JH, Kim JC, Yu CS, Park SH, et al. Incremental value

of liver MR imaging in patients with potentially curable colorectal hepatic metastasis detected at CT: A prospective comparison of diffusion-weighted imaging, gadoxetic acid-enhanced MR imaging, and a combination of both MR techniques. *Radiology*. 2015;**274**(3):712-722

[73] Seale MK, Catalano OA, Saini S, Hahn PF, Sahani DV. Hepatobiliary-specific MR contrast agents: Role in imaging the liver and biliary tree. *Radiographics*. 2009;**29**(6):1725-1748

[74] Koh DM, Berry J. Critical questions in the imaging of colorectal hepatic metastases. *Cancer Imaging*. 2008;**8** (Spec Iss A):S69-S78

[75] Hindman JC. Proton resonance shift of water in the gas and liquid states. *The Journal of Chemical Physics*. 1966;**44**(12):4582-4592

[76] Ishihara Y, Calderon A, Watanabe H, Okamoto K, Suzuki Y, Kuroda K, et al. A precise and fast temperature mapping using water proton chemical shift. *Magnetic Resonance in Medicine*. 1995;**34**(6):814-823

[77] Poorter JD, Wagter CD, Deene YD, Thomsen C, Stahlberg F, Achten E. Noninvasive MRI thermometry with the proton resonance frequency (PRF) method: In vivo results in human muscle. *Magnetic Resonance in Medicine*. 1995;**33**(1):74-81

[78] Rieke V, Butts PK. MR thermometry. *Journal of Magnetic Resonance Imaging*. 2008;**27**(2):376-390

[79] Quesson B, de Zwart JA, Moonen CTW. Magnetic resonance temperature imaging for guidance of thermotherapy. *Journal of Magnetic Resonance Imaging*. 2000;**12**(4):525-533

- [80] Haase A, Frahm J, Matthaei D, Hanicke W, Merboldt KD. FLASH imaging: Rapid NMR imaging using low flip-angle pulses. 1986. *Journal of Magnetic Resonance*. 2011;**213**(2):533-541
- [81] Auboiroux V, Viallon M, Roland J, Hyacinthe J, Petrusca L, Morel DR, et al. ARFI-prepared MRgHIFU in liver: Simultaneous mapping of ARFI-displacement and temperature elevation, using a fast GRE-EPI sequence. *Magnetic Resonance in Medicine*. 2012;**68**(3):932-946
- [82] McDannold N, Maier SE. Magnetic resonance acoustic radiation force imaging. *Medical Physics*. 2008;**35**(8):3748-3758
- [83] Payne A, Merrill R, Minalga E, Hadley JR, Odeen H, Hofstetter LW, et al. A breast-specific MR guided focused ultrasound platform and treatment protocol: First-in-human technical evaluation. *IEEE Transactions on Biomedical Engineering*. 2021;**68**(3):893-904
- [84] Bour P, Marquet F, Ozenne V, Toupin S, Dumont E, Aubry JF, et al. Real-time monitoring of tissue displacement and temperature changes during MR-guided high intensity focused ultrasound. *Magnetic Resonance in Medicine*. 2017;**78**(5):1911-1921
- [85] Paquin R, Vignaud A, Marsac L, Younan Y, Lehericy S, Tanter M, et al. Keyhole acceleration for magnetic resonance acoustic radiation force imaging (MR ARFI). *Magnetic Resonance Imaging*. 2013;**31**(10):1695-1703
- [86] Kaye EA, Hertzberg Y, Marx M, Werner B, Navon G, Levoy M, et al. Application of Zernike polynomials towards accelerated adaptive focusing of transcranial high intensity focused ultrasound. *Medical Physics*. 2012;**39**(10):6254-6263
- [87] Hertzberg Y, Volovick A, Zur Y, Medan Y, Vitek S, Navon G. Ultrasound focusing using magnetic resonance acoustic radiation force imaging: Application to ultrasound transcranial therapy. *Medical Physics*. 2010;**37**(6 Part 1):2934-2942
- [88] Mansfield P. Snapshot magnetic resonance imaging (Nobel lecture). *Angewandte Chemie (International Ed. in English)*. 2004;**43**(41):5456-5464
- [89] Stehling MK, Turner R, Mansfield P. Echo-planar imaging: Magnetic resonance imaging in a fraction of a second. *Science*. 1991;**254**(5028):43-50
- [90] Zhang S, Block KT, Frahm J. Magnetic resonance imaging in real time: Advances using radial FLASH. *Journal of Magnetic Resonance Imaging*. 2010;**31**(1):101-109
- [91] McIntyre D, Bowtell R, Commandre M, Hennel F, Mansfield P, Morris P. Overcoming field inhomogeneities in MRI. In: *The Bio-Clinical Interface: Bio-Clinical Psychiatry Mapping Brain Function*; Edited Proceedings of the Bio-Clinical Interface Conferences, Held in Rouffach, France, between 1992 and 1994. Arcueil, France: John Libbey Eurotext. pp. 349-355
- [92] Salomir R, de Senneville BD, Moonen CT. A fast calculation method for magnetic field inhomogeneity due to an arbitrary distribution of bulk susceptibility. *Concepts in Magnetic Resonance Part B: Magnetic Resonance Engineering*. 2003;**19**(1):26-34
- [93] Hayashi M, Inoue Y, Komeda K, Shimizu T, Asakuma M, Hirokawa F,



- et al. Clinicopathological analysis of recurrence patterns and prognostic factors for survival after hepatectomy for colorectal liver metastasis. *BMC Surgery*. 2010;**10**(1):27
- [94] Yamashita S, Chun YS, Kopetz SE, Vauthey JN. Biomarkers in colorectal liver metastases. *The British Journal of Surgery*. 2018;**105**(6):618-627
- [95] Adam R, De Gramont A, Figueras J, Guthrie A, Kokudo N, Kunstlinger F, et al. The oncosurgery approach to managing liver metastases from colorectal cancer: A multidisciplinary international consensus. *The Oncologist*. 2012;**17**(10):1225-1239
- [96] Blumgart LH, Leach KG, Karran SJ. Observations on liver regeneration after right hepatic lobectomy. *Gut*. 1971;**12**(11):922-928
- [97] Fang C-H, Liu J, Fang Fan Y, Yang J, Xiang N, Zeng N. Outcomes of hepatectomy for hepatolithiasis based on 3-dimensional reconstruction technique. *Journal of the American College of Surgeons*. 2013;**217**(2):280-288
- [98] Majno P, Azoulay D, Bismuth H. In: Bircher J, Benhamou JP, McIntyre N, Rizzetto M, Rodes J, editors. *General Surgical Aspects and the Risks of Liver Surgery in Patients with Hepatic Disease*. Oxford Textbook of Clinical Hepatology. Oxford: Oxford University Press; 1999. pp. 1997-2008
- [99] Aloia TA, Vauthey JN. Chapter 92—Hepatic metastasis from colorectal cancer. In: Jarnagin WR, editor. *Blumgart's Surgery of the Liver, Biliary Tract and Pancreas, 2-Volume Set (Sixth Edition)*. 6th ed. Philadelphia: Elsevier; 2017. pp. 1339-54.e4
- [100] Mentha G, Roth AD, Terraz S, Giostra E, Gervaz P, Andres A, et al. 'Liver first' approach in the treatment of colorectal cancer with synchronous liver metastases. *Digestive Surgery*. 2008;**25**(6):430-435
- [101] Jegatheeswaran S, Mason JM, Hancock HC, Siriwardena AK. The liver-first approach to the management of colorectal cancer with synchronous hepatic metastases: A systematic review. *JAMA Surgery*. 2013;**148**(4):385-391
- [102] Berber E, Pelley R, Siperstein AE. Predictors of survival after radiofrequency thermal ablation of colorectal cancer metastases to the liver: A prospective study. *Journal of Clinical Oncology*. 2005;**23**(7):1358-1364
- [103] Lau WY. *Different Approaches to Liver Resection*. Singapore: Springer Singapore; 2021. pp. 155-169
- [104] Lai ECH, Lau SHY, Lau WY. In: Yan L, editor. *Anatomical Liver Resection*. Dordrecht: Springer Netherlands; 2016. pp. 111-116
- [105] Lortat-Jacob J, Robert H. Well defined technic for right hepatectomy. *La Presse Médicale*. 1952;**60**(26):549-551
- [106] Pack GT, Baker HW. Total right hepatic lobectomy: Report of a case. *Annals of Surgery*. 1953;**138**(2):253
- [107] Woodington GF, Waugh JM. Results of resection of metastatic tumors of the liver. *American Journal of Surgery*. 1963;**105**(1):24-29
- [108] Adam R, Laurent A, Azoulay D, Castaing D, Bismuth H. Two-stage hepatectomy: A planned strategy to treat irresectable liver tumors. *Annals of Surgery*. 2000;**232**(6):777-785
- [109] Alvarez FA, Ardiles V, Sanchez Claria R, Pekolj J, de Santibanes E. Associating liver partition and portal

- vein ligation for staged hepatectomy (ALPPS): Tips and tricks. *Journal of Gastrointestinal Surgery*. 2013;**17**(4): 814-821
- [110] de Santibanes E, Clavien PA. Playing play-doh to prevent postoperative liver failure: The "ALPPS" approach. *Annals of Surgery*. 2012;**255**(3):415-417
- [111] Makuuchi M, Thai BL, Takayasu K, Takayama T, Kosuge T, Gunven P, et al. Preoperative portal embolization to increase safety of major hepatectomy for hilar bile duct carcinoma: A preliminary report. *Surgery*. 1990;**107**(5):521-527
- [112] Jaeck D, Oussoultzoglou E, Rosso E, Greget M, Weber JC, Bachellier P. A two-stage hepatectomy procedure combined with portal vein embolization to achieve curative resection for initially unresectable multiple and bilobar colorectal liver metastases. *Annals of Surgery*. 2004;**240**(6):1037-1049; discussion 1049-51
- [113] Schadde E, Clavien PA. In: Clavien PA, Sarr MG, Fong Y, Miyazaki M, editors. *ALPPS (Associating Liver Partition with Portal Vein Ligation for Staged Hepatectomy)*. Berlin, Heidelberg: Springer Berlin Heidelberg; 2016. pp. 447-456
- [114] Dueland S, Syversveen T, Solheim JM, Solberg S, Grut H, Bjørneth BA, et al. Survival following liver transplantation for patients with nonresectable liver-only colorectal metastases. *Annals of Surgery*. 2020;**271**(2):212-218
- [115] Andres A, Toso C. Liver transplantation for colorectal metastasis: A stepwise approach. *Hepatobiliary Surgery and Nutrition*. 2020;**9**(3):336
- [116] Torres-Jimenez J, Esteban-Villarrubia J, Ferreiro-Monteagudo R, Carrato A. Local Treatments in the unresectable patient with colorectal cancer metastasis: A review from the point of view of the medical oncologist. *Cancers*. 2021;**13**(23)
- [117] Gallinger S, Biagi JJ, Fletcher GG, Nhan C, Ruo L, McLeod RS. Liver resection for colorectal cancer metastases. *Current Oncology*. 2013;**20**(3):e255-e265
- [118] Lam VW, Spiro C, Laurence JM, Johnston E, Hollands MJ, Pleass HC, et al. A systematic review of clinical response and survival outcomes of downsizing systemic chemotherapy and rescue liver surgery in patients with initially unresectable colorectal liver metastases. *Annals of Surgical Oncology*. 2012;**19**(4):1292-1301
- [119] Mentha G, Majno P, Terraz S, Rubbia-Brandt L, Gervaz P, Andres A, et al. Treatment strategies for the management of advanced colorectal liver metastases detected synchronously with the primary tumour. *European Journal of Surgical Oncology*. 2007;**33**(Suppl. 2):S76-S83
- [120] Khoo E, O'Neill S, Brown E, Wigmore SJ, Harrison EM. Systematic review of systemic adjuvant, neoadjuvant and perioperative chemotherapy for resectable colorectal-liver metastases. *HPB: The Official Journal of the International Hepato Pancreato Biliary Association*. 2016;**18**(6):485-493
- [121] Bierman HR, Byron JRL, Kelley KH, Grady A. Studies on the blood supply of tumors in man. III. Vascular patterns of the liver by hepatic arteriography in vivo. *Journal of the National Cancer Institute*. 1951;**12**(1):107-131
- [122] Breedis C, Young G. The blood supply of neoplasms in the liver. *The American Journal of Pathology*. 1954;**30**(5):969-977

- [123] Ranieri G, Laforgia M, Nardulli P, Ferraiuolo S, Molinari P, Marech I, et al. Oxaliplatin-based intra-arterial chemotherapy in colo-rectal cancer liver metastases: A review from pharmacology to clinical application. *Cancers (Basel)*. 2019;**11**(2):141-156
- [124] Minchinton AI, Tannock IF. Drug penetration in solid tumours. *Nature Reviews. Cancer*. 2006;**6**(8):583-592
- [125] Kemeny MM, Battifora H, Blayney DW, Cecchi G, Goldberg DA, Leong LA, et al. Sclerosing cholangitis after continuous hepatic artery infusion of FUDR. *Annals of Surgery*. 1985;**202**(2):176-181
- [126] Beltrañ-Gracia E, Lopez-Camacho A, Higuera-Ciapa I, Velazquez-Fernandez JB, Vallejo-Cardona AA. Nanomedicine review: Clinical developments in liposomal applications. *Cancer Nanotechnology*. 2019;**10**(1):1-40
- [127] Lyon PC, Griffiths LF, Lee J, Chung D, Carlisle R, Wu F, et al. Clinical trial protocol for TARDOX: A phase I study to investigate the feasibility of targeted release of lyso-thermosensitive liposomal doxorubicin (ThermoDox®) using focused ultrasound in patients with liver tumours. *Journal of Therapeutic Ultrasound*. 2017;**5**(1):1-8
- [128] Gray MD, Lyon PC, Mannaris C, Folkes LK, Stratford M, Campo L, et al. Focused ultrasound hyperthermia for targeted drug release from thermosensitive liposomes: Results from a Phase I Trial. *Radiology*. 2019;**291**(1):232-238
- [129] National Library of Medicine (U.S.). A Phase I Study of Lyso-thermosensitive Liposomal Doxorubicin and MR-HIFU for Pediatric Refractory Solid Tumors. Identifier NCT02536183. 2015. Available from: <https://clinicaltrials.gov/ct2/show/NCT02536183>
- [130] National Library of Medicine (U.S.) Ultrasound-enhanced Delivery of Chemotherapy to Patients With Liver Metastasis From Breast- and Colorectal Cancer. Identifier NCT03477019. 2018. Available from: <https://clinicaltrials.gov/ct2/show/NCT03477019>
- [131] Cheng JC, Wu JK, Huang CM, Liu HS, Huang DY, Cheng SH, et al. Radiation-induced liver disease after three-dimensional conformal radiotherapy for patients with hepatocellular carcinoma: Dosimetric analysis and implication. *International Journal of Radiation Oncology, Biology, Physics*. 2002;**54**(1):156-162
- [132] Pan CC, Kavanagh BD, Dawson LA, Li XA, Das SK, Miften M, et al. Radiation-associated liver injury. *International Journal of Radiation Oncology, Biology, Physics*. 2010;**76**(3 Suppl):S94-S100
- [133] Kirichenko A, Gayou O, Parda D, Kudithipudi V, Tom K, Khan A, et al. Stereotactic body radiotherapy (SBRT) with or without surgery for primary and metastatic liver tumors. *HPB*. 2016;**18**(1):88-97
- [134] Ursino S, Greco C, Cartei F, Colosimo C, Stefanelli A, Cacopardo B, et al. Radiotherapy and hepatocellular carcinoma: Update and review of the literature. *European Review for Medical and Pharmacological Sciences*. 2012;**16**(11):1599-1604
- [135] Comito T, Clerici E, Tozzi A, D'Agostino G. Liver metastases and SBRT: A new paradigm? *Reports of Practical Oncology and Radiotherapy*. 2015;**20**(6):464-471
- [136] Aitken KL, Hawkins MA. Stereotactic body radiotherapy for liver metastases. *Clinical Oncology (Royal College of Radiologists)*. 2015;**27**(5):307-315

- [137] Petrelli F, Comito T, Barni S, Pancera G, Scorsetti M, Ghidini A. Stereotactic body radiotherapy for colorectal cancer liver metastases: A systematic review. *Radiotherapy and Oncology*. 2018;**129**(3):427-434
- [138] National Library of Medicine (U.S.). The MOMENTUM Study: The Multiple Outcome Evaluation of Radiation Therapy Using the MR-Linac Study (MOMENTUM). Identifier NCT04075305. 2019. Available from: <https://clinicaltrials.gov/ct2/show/NCT04075305>
- [139] National Library of Medicine (U.S.). Feasibility of Online MR-guided Radiotherapy on a 1.5T MR-Linac. Identifier NCT04172753. 2019. Available from: <https://clinicaltrials.gov/ct2/show/NCT04172753>
- [140] National Library of Medicine (U.S.). OAR-Based, Dose Escalated SBRT With Real Time Adaptive MRI Guidance for Liver Metastases. Identifier NCT04020276. 2019. Available from: <https://clinicaltrials.gov/ct2/show/NCT04020276>
- [141] National Library of Medicine (U.S.). Phase II of Adaptive Magnetic Resonance-Guided Stereotactic Body Radiotherapy (SBRT) for Treatment of Primary or Secondary Progressive Liver Tumors. Identifier NCT04242342. 2019. Available from: <https://clinicaltrials.gov/ct2/show/NCT04242342>
- [142] Czito B. Magnetic Resonance Imaging (MRI) Solely For Liver Stereotactic Body Radiation Therapy (SBRT). Identifier NCT02722395. 2016-2017. Available from: <https://clinicaltrials.gov/ct2/show/NCT02722395>
- [143] National Library of Medicine (U.S.). MRI-Guided Stereotactic Body Radiation Therapy in Treating Patients With Liver Metastases or Liver Cancer. Identifier NCT02683200. 2016. Available from: <https://clinicaltrials.gov/ct2/show/NCT02683200>
- [144] van Dams R, Wu TC, Kishan AU, Raldow AC, Chu FI, Hernandez J, et al. Ablative radiotherapy for liver tumors using stereotactic MRI-guidance: A prospective phase I trial. *Radiotherapy and Oncology*. 2022;**170**:14-20
- [145] Giles SL, Brown MRD, Rivens I, Deppe M, Huisman M, Kim YS, et al. Comparison of imaging changes and pain responses in patients with intra- or extraosseous bone metastases treated palliatively with magnetic resonance-guided high-intensity-focused ultrasound. *Journal of Vascular and Interventional Radiology*. 2019;**30**(9):1351-60 e1
- [146] Liberman B, Gianfelice D, Inbar Y, Beck A, Rabin T, Shabshin N, et al. Pain palliation in patients with bone metastases using MR-guided focused ultrasound surgery: A multicenter study. *Annals of Surgical Oncology*. 2009;**16**(1):140-146
- [147] AbdElmagied AM, Vaughan LE, Weaver AL, Laughlin-Tommaso SK, Hesley GK, Woodrum DA, et al. Fibroid interventions: Reducing symptoms today and tomorrow: Extending generalizability by using a comprehensive cohort design with a randomized controlled trial. *American Journal of Obstetrics and Gynecology*. 2016;**215**(3):338 e1-338 e18
- [148] Chen R, Keserci B, Bi H, Han X, Wang X, Bai W, et al. The safety and effectiveness of volumetric magnetic resonance-guided high-intensity focused ultrasound treatment of symptomatic uterine fibroids: Early clinical experience in China. *Journal of Therapeutic Ultrasound*. 2016;**4**(1):27

- [149] Dorenberg EJ, Courivaud F, Ring E, Hald K, Jakobsen JA, Fosse E, et al. Volumetric ablation of uterine fibroids using Sonalleve high-intensity focused ultrasound in a 3 Tesla scanner—first clinical assessment. *Minimally Invasive Therapy & Allied Technologies*. 2013;**22**(2):73-79
- [150] Laughlin-Tommaso S, Barnard EP, AbdElmagied AM, Vaughan LE, Weaver AL, Hesley GK, et al. FIRSST study: Randomized controlled trial of uterine artery embolization vs focused ultrasound surgery. *American Journal of Obstetrics and Gynecology*. 2019;**220**(2):174 e1-174 e13
- [151] Verpalen IM, de Boer JP, Linstra M, Pol RLI, Nijholt IM, Moonen CTW, et al. The Focused Ultrasound Myoma Outcome Study (FUMOS); a retrospective cohort study on long-term outcomes of MR-HIFU therapy. *European Radiology*. 2020;**30**(5):2473-2482
- [152] Giles SL, Imseeh G, Rivens I, Ter Haar GR, Taylor A, deSouza NM. MR guided high intensity focused ultrasound (MRgHIFU) for treating recurrent gynaecological tumours: A pilot feasibility study. *The British Journal of Radiology*. 2019;**92**(1098):20181037
- [153] Napoli A, Anzidei M, De Nunzio C, Cartocci G, Panebianco V, De Dominicis C, et al. Real-time magnetic resonance-guided high-intensity focused ultrasound focal therapy for localised prostate cancer: Preliminary experience. *European Urology*. 2013;**63**(2):395-398
- [154] Tay KJ, Cheng CWS, Lau WKO, Khoo J, Thng CH, Kwek JW. Focal therapy for prostate cancer with in-BORE MR-guided focused ultrasound: Two-year follow-up of a phase I trial—complications and functional outcomes. *Radiology*. 2017;**285**(2):620-628
- [155] Chang JW, Park CK, Lipsman N, Schwartz ML, Ghanouni P, Henderson JM, et al. A prospective trial of magnetic resonance-guided focused ultrasound thalamotomy for essential tremor: Results at the 2-year follow-up. *Annals of Neurology*. 2018;**83**(1):107-114
- [156] Elias WJ, Lipsman N, Ondo WG, Ghanouni P, Kim YG, Lee W, et al. A randomized trial of focused ultrasound thalamotomy for essential tremor. *The New England Journal of Medicine*. 2016;**375**(8):730-739
- [157] Furusawa H, Namba K, Thomsen S, Akiyama F, Bendet A, Tanaka C, et al. Magnetic resonance-guided focused ultrasound surgery of breast cancer: Reliability and effectiveness. *Journal of the American College of Surgeons*. 2006;**203**(1):54-63
- [158] Wu F, Chen WZ, Bai J, Zou JZ, Wang ZL, Zhu H, et al. Pathological changes in human malignant carcinoma treated with high-intensity focused ultrasound. *Ultrasound in Medicine & Biology*. 2001;**27**(8):1099-1106
- [159] Wu F, Wang ZB, Chen WZ, Wang W, Gui Y, Zhang M, et al. Extracorporeal high intensity focused ultrasound ablation in the treatment of 1038 patients with solid carcinomas in China: An overview. *Ultrasonics Sonochemistry*. 2004;**11**(3):-149, 54. 4th Conference on the Applications of Power Ultrasound in Physical and Chemical Processing
- [160] Wu F, Wang ZB, Chen WZ, Zhu H, Bai J, Zou JZ, et al. Extracorporeal high intensity focused ultrasound ablation in the treatment of patients with large hepatocellular carcinoma. *Annals of Surgical Oncology*. 2004;**11**(12):1061-1069
- [161] Yang T, Ng DM, Du N, He N, Dai X, Chen P, et al. HIFU for the treatment

of difficult colorectal liver metastases with unsuitable indications for resection and radiofrequency ablation: A phase I clinical trial. *Surgical Endoscopy*. 2021;**35**(5):2306-2315

[162] Tsang SH, Ma KW, She WH, Chu F, Lau V, Lam SW, et al. High-intensity focused ultrasound ablation of liver tumors in difficult locations. *International Journal of Hyperthermia*. 2021;**38**(2):56-64

[163] Park MY, Jung SE, Cho SH, Piao XH, Hahn ST, Han JY, et al. Preliminary experience using high intensity focused ultrasound for treating liver metastasis from colon and stomach cancer. *International Journal of Hyperthermia*. 2009;**25**(3):180-188

[164] Illing R, Kennedy J, Wu F, Ter Haar G, Protheroe A, Friend P, et al. The safety and feasibility of extracorporeal high-intensity focused ultrasound (HIFU) for the treatment of liver and kidney tumours in a Western population. *British Journal of Cancer*. 2005;**93**(8):890-895

[165] Orsi F, Zhang L, Arnone P, Orgera G, Bonomo G, Vigna PD, et al. High-intensity focused ultrasound ablation: Effective and safe therapy for solid tumors in difficult locations. *AJR. American Journal of Roentgenology*. 2010;**195**(3):W245-W252

[166] Sehmbi AS, Froghi S, Oliveira de Andrade M, Saffari N, Fuller B, Quaglia A, et al. Systematic review of the role of high intensity focused ultrasound (HIFU) in treating malignant lesions of the hepatobiliary system. *HPB*. 2021;**23**(2):187-196

[167] Schuster TG, Wei JT, Hendlin K, Jahnke R, Roberts WW. Histotripsy treatment of benign prostatic enlargement using the Vortx Rx system:

Initial human safety and efficacy outcomes. *Urology*. 2018;**114**:184-187

[168] Solbiati LA. The HistoSonics System for Treatment of Primary and Metastatic Liver Tumors Using Histotripsy (#HOPE4LIVER). Identifier NCT04573881. 2020. Available from: <https://clinicaltrials.gov/ct2/show/NCT04573881>

[169] Cranston D. A review of high intensity focused ultrasound in relation to the treatment of renal tumours and other malignancies. *Ultrasonics Sonochemistry*. 2015;**27**:654-658

[170] Gedroyc WM. New clinical applications of magnetic resonance-guided focused ultrasound. *Topics in Magnetic Resonance Imaging*. 2006;**17**(3):189-194

[171] Diana M, Schiraldi L, Liu YY, Memeo R, Mutter D, Pessaux P, et al. High intensity focused ultrasound (HIFU) applied to hepato-bilio-pancreatic and the digestive system-current state of the art and future perspectives. *Hepatobiliary Surgery and Nutrition*. 2016;**5**(4):329-344

[172] Desgranges S, Lorton O, Gui-Levy L, Guillemin P, Celicanin Z, Hyacinthe JN, et al. Micron-sized PFOB liquid core droplets stabilized with tailored-made perfluorinated surfactants as a new class of endovascular sono-sensitizers for focused ultrasound thermotherapy. *Journal of Materials Chemistry B*. 2019;**7**(6):927-939

[173] Guillemin PC, Gui L, Lorton O, Zilli T, Crowe LA, Desgranges S, et al. Mild hyperthermia by MR-guided focused ultrasound in an ex vivo model of osteolytic bone tumour: Optimization of the spatio-temporal control of the delivered temperature. *Journal of Translational Medicine*. 2019;**17**(1):350

- [174] Mori N, Jud C, Salomir R, Cattin PC. Leveraging respiratory organ motion for non-invasive tumor treatment devices: A feasibility study. *Physics in Medicine and Biology*. 2016;**61**(11):4247-4267
- [175] Muller A, Petrusca L, Auboiroux V, Valette PJ, Salomir R, Cotton F. Management of respiratory motion in extracorporeal high-intensity focused ultrasound treatment in upper abdominal organs: Current status and perspectives. *Cardiovascular and Interventional Radiology*. 2013;**36**(6):1464-1476
- [176] Salomir R, Petrusca L, Auboiroux V, Muller A, Vargas MI, Morel DR, et al. Magnetic resonance-guided shielding of prefocal acoustic obstacles in focused ultrasound therapy: Application to intercostal ablation in liver. *Investigative Radiology*. 2013;**48**(6):366-380
- [177] Jung SE, Cho SH, Jang JH, Han JY. High-intensity focused ultrasound ablation in hepatic and pancreatic cancer: Complications. *Abdominal Imaging*. 2011;**36**(2):185-195
- [178] Zhang L, Zhu H, Jin C, Zhou K, Li K, Su H, et al. High-intensity focused ultrasound (HIFU): Effective and safe therapy for hepatocellular carcinoma adjacent to major hepatic veins. *European Radiology*. 2009;**19**(2):437-445
- [179] Cheung TT, Poon RTP, Yau T, Tsang DSF, Lo CM, Fan ST. High-intensity focused ultrasound as a treatment for colorectal liver metastasis in difficult position. *International Journal of Colorectal Disease*. 2012;**27**(7):987-988
- [180] Preiswerk F, De Luca V, Arnold P, Celicanin Z, Petrusca L, Tanner C, et al. Model-guided respiratory organ motion prediction of the liver from 2D ultrasound. *Medical Image Analysis*. 2014;**18**(5):740-751
- [181] de Zwart JA, Vimeux FC, Palussiere J, Salomir R, Quesson B, Delalande C, et al. On-line correction and visualization of motion during MRI-controlled hyperthermia. *Magnetic Resonance in Medicine*. 2001;**45**(1):128-137
- [182] Lorton O, Guillemain PC, M'Rad Y, Peloso A, Boudabbous S, Charbonnier C, et al. A novel concept of a phased-array HIFU transducer optimized for MR-guided hepatic ablation: Embodiment and first in-vivo studies. *Frontiers in Oncology*. 2022;**12**
- [183] Lorton O, Guillemain PC, Holman R, Desgranges S, Gui L, Crowe LA, et al. Enhancement of HIFU thermal therapy in perfused tissue models using micron-sized FTAC-stabilized PFOB-core endovascular sonosensitizers [Journal Article]. *International Journal of Hyperthermia*. 2020;**37**(1):1116-1130
- [184] Astafyeva K, Somaglino L, Desgranges S, Berti R, Patinote C, Langevin D, et al. Perfluorocarbon nanodroplets stabilized by fluorinated surfactants: Characterization and potentiality as theranostic agents. *Journal of Materials Chemistry B*. 2015;**3**(14):2892-2907
- [185] Lorton O, Guillemain PC, Holman R, Desgranges S, Gui L, Crowe LA, et al. Enhancement of HIFU thermal therapy in perfused tissue models using micron-sized FTAC-stabilized PFOB-core endovascular sonosensitizers. *International Journal of Hyperthermia*. 2020;**37**(1):1116-1130
- [186] Holman R, Gui L, Lorton O, Guillemain P, Desgranges S, Contino-Pépin C, et al. PFOB sonosensitive microdroplets: Determining their interaction radii with focused ultrasound using MR thermometry and a Gaussian convolution kernel computation.

International Journal of Hyperthermia. 2022;**39**(1):108-119

[187] Okada A, Murakami T, Mikami K, Onishi H, Tanigawa N, Marukawa T, et al. A case of hepatocellular carcinoma treated by MR-guided focused ultrasound ablation with respiratory gating. *Magnetic Resonance in Medical Sciences*. 2006;**5**(3):167-171

[188] Anzidei M, Napoli A, Sandolo F, Marincola BC, Di Martino M, Berloco P, et al. Magnetic resonance-guided focused ultrasound ablation in abdominal moving organs: A feasibility study in selected cases of pancreatic and liver cancer. *Cardiovascular and Interventional Radiology*. 2014;**37**(6):1611-1617

[189] Gedroyc WM. MR guided focused ultrasound of the liver. In: 2nd International MR-guided Focused Ultrasound Symposium, Focused Ultrasound Foundation. 2010. Available from: <https://www.fusfoundation.org>

[190] Dupré A, Melodelima D, Perol D, Chen Y, Vincenot J, Chapelon JY, et al. First clinical experience of intra-operative high intensity focused ultrasound in patients with colorectal liver metastases: A phase I-IIa study. *PLoS One*. 2015;**10**(2):e0118212

[191] Dupré A, Perol D, Blanc E, Peyrat P, Basso V, Chen Y, et al. Efficacy of high-intensity focused ultrasound-assisted hepatic resection (HIFU-AR) on blood loss reduction in patients with liver metastases requiring hepatectomy: Study protocol for a randomized controlled trial. *Trials*. 2017;**18**(1):1-10

[192] Cambronero S, Rivoire M, Dupré A, Melodelima D. Non-invasive fast, large and selective in vivo HIFU ablation of the liver with a toroidal transducer. In: 2019 IEEE International Ultrasonics Symposium (IUS). 2019. pp. 2126-2128

[193] Dupre A, Melodelima D, Perol D, Chen Y, Vincenot J, Chapelon JY, et al. Evaluation of the feasibility, safety, and accuracy of an intraoperative high-intensity focused ultrasound device for treating liver metastases. *Journal of Visualized Experiments*. 2019;**143**:e57964

[194] Vaezy S, Zderic V. Hemorrhage control using high intensity focused ultrasound. *International Journal of Hyperthermia*. 2007;**23**(2):203-211

[195] Silveira LA, Silveira FBC, Fazan VPS. Arterial diameter of the celiac trunk and its branches: Anatomical study. *Acta Cirúrgica Brasileira*. 2009;**24**(1):43-47

[196] Zaccagna F, Anzidei M, Sandolo F, Cavallo Marincola B, Palla C, Leonardi A, et al. MRgFUS for liver and pancreas cancer treatments: The Umberto I hospital experience. *Translational Cancer Research*. 2014;**3**(5):430-441

[197] Fischer K, Gedroyc W, Jolesz FA. Focused ultrasound as a local therapy for liver cancer. *Cancer Journal*. 2010;**16**(2):118-124

[198] National Library of Medicine (U.S.). MR-guided High Intensity Focused Ultrasound (HIFU) on Pediatric Solid Tumors. Identifier NCT02076906. 2014. Available from: <https://clinicaltrials.gov/ct2/show/NCT02076906>

[199] Zhang Q, Bian S, Lv W, Kou D, Hu H, Guo S, et al. Observation of efficacy of TACE combined with HIFU on patients with middle-advanced liver cancer. *European Review for Medical and Pharmacological Sciences*. 2019;**23**(3):239-246

[200] Luo Y, Jiang Y. Comparison of efficiency of TACE plus HIFU and TACE alone on patients with primary liver cancer. *Journal of the College of*



Physicians and Surgeons–Pakistan.  
2019;**29**(5):414-417

[201] Wijlemans JW, Bartels LW,  
Deckers R, Ries M, Mali WP, Moonen CT,  
et al. Magnetic resonance-guided high-  
intensity focused ultrasound (MR-HIFU)  
ablation of liver tumours. *Cancer  
Imaging*. 2012;**12**(2):387-394

IntechOpen

IntechOpen